

Lattice construction of 't Hooft anomaly with \mathbb{Z}_N 1-form symmetry

Motokazu Abe (Kyushu U.)

Seminar @ Osaka U.

2023/11/21

Topology of SU(N) lattice gauge theories coupled with \mathbb{Z}_N 2-form gauge fields

M. Abe, O. Morikawa, S. Onoda, H. Suzuki and Y. Tanizaki

arXiv:2303.10977[hep-th]

Fractional topological charge in lattice Abelian gauge theory

M. Abe, O. Morikawa and H. Suzuki

PTEP 2023 (2023) 2, 023B03 [arXiv:2210.12967[hep-th]]

Symmetry and Anomaly I

- Classical Theory : **Symmetry** \longleftrightarrow **Conservation law** (Noether Theorem)
- Quantum Theory : The conservation law may be broken (Anomaly).
 - Focus on **the Partition function**,

$$Z[A] = \int [\mathcal{D}(\text{field})] e^{S[\text{field}, A]}.$$

- How to distinguish the anomaly : Whether the Z is **invariant or not** under a transformation

$$Z' \stackrel{?}{=} Z$$

$$\begin{aligned} \rightarrow Z'[A + \partial\theta] &= \int [\mathcal{D}(\text{field})] e^{S[\text{field}, A + \partial\theta]} \\ &= \underbrace{e^{\mathcal{A}[\theta, A]}}_{\text{'t Hooft anomaly}} \underbrace{\int [\mathcal{D}(\text{field})] e^{S[\text{field}, A]}}_{= Z}. \end{aligned}$$

Symmetry and Anomaly II


- We can predict the **low energy dynamics** of the gauge theory.
 - ✂ Gauge theory : A theory which describes the Standard Model of particles
- ✓ e. g., we decided the theory for the strong interaction is the SU(3) gauge theory because **the theory** and **the experiment** are well matched.

Particle Theory



Predict

Particle Experiment



<https://www.icepp.s.u-tokyo.ac.jp/information/20220426.html>

High

Low


Energy

Recent Developments in Anomalies

- Recently, Gaiotto et al. has extended the concept of symmetry. : Higher Form Symmetry (Gaiotto, Kapustin, Seiberg, Willett, arXiv:1412.5148[hep-th])
 - By anomalies with higher form (and discrete) symmetries, the low energy dynamics of gauge theories has been discussed. (Gaiotto, Kapustin, Komargodski, Seiberg, arXiv:1703.00501[hep-th])
 - Many new anomalies have been discovered and related studies has been done.
 - ✓ Yamaguchi, arXiv:1811.09390[hep-th]
 - ✓ Hidaka, Hirono, Nitta, Tanizaki, Yokokura, arXiv:1903.06389[hep-th]
 - ✓ Honda, Tanizaki, arXiv:2009.10183[hep-th]
 - ✓ etc.

☆Motivation : Understand these anomalies in the **lattice field theory** where we treat the regularization well.

Lattice Gauge Theory



Higher Form Symmetry I


Gaiotto, Kapustin, Seiberg, Willett, arXiv:1412.5148[hep-th]

- Traditional symmetry (0-form symmetry) : transform the point $\psi(x)$.

✓ e.g., global $U(1)$ symmetry $\psi(x) \rightarrow e^{i\alpha}\psi(x)$.

- In (2+1)d, look this $\psi(x)$'s transformation by the symmetry operator,

$$e^{i\alpha\Omega}\psi(x)e^{-i\alpha\Omega} = e^{-i\alpha}\psi(x), \quad \Omega = \int_M d^2x j^0(x), \quad j^\mu(x) = i\bar{\psi}(x)\gamma^\mu\psi(x).$$




Higher Form Symmetry II

Gaiotto, Kapustin, Seiberg, Willett, arXiv:1412.5148[hep-th]

- Traditional symmetry (0-form symmetry) : transform the point $\psi(x)$.
 - ✓ e.g., global $U(1)$ symmetry $\psi(x) \rightarrow e^{i\alpha}\psi(x)$.
- Extend the point to 2d, 3d, ... objects


- 0-form symmetry

- Transform **point** $\psi(x)$



- 1-form symmetry

- Transform **loop** $W(C)$



Anomaly of the $SU(N)$ gauge theory with θ term

- The $SU(N)$ gauge theory with the θ term has the time reversal (\mathcal{T}) symmetry at $\theta = \pi$.

$$Z = \int \mathcal{D}a e^{S[a]} = \int \mathcal{D}a e^{S_{SU(N)}[a]} e^{i\theta Q[a]}, \quad Q \in \mathbb{Z}$$

$$\xrightarrow{\theta=\pi, \mathcal{T} \text{ trans.}} Z' = \int \mathcal{D}a e^{S_{SU(N)}[a]} e^{i\pi(-Q[a])} = \int \mathcal{D}a e^{S_{SU(N)}[a]} e^{i\pi(+Q[a])} \underbrace{e^{-i2\pi Q[a]}}_{=1} = Z$$

- Then, we construct the $SU(N)$ gauge theory with the higher form symmetry (\mathbb{Z}_N 1-form gauge symmetry). This means we couple \mathbb{Z}_N 2-form gauge field to the theory.
 - The topological charge (TC) becomes fractional, so it is not invariant under the \mathcal{T} transformation.

Important!!

$$e^{-i2\pi Q} \neq 1$$

- This theory at $\theta = \pi$ has the mixed anomaly between the \mathbb{Z}_N 1-form gauge and \mathcal{T} symmetry.

Topological Charge on the Lattice

- How to calculate the topological charge Q ,


$$Q = -\frac{1}{24\pi^2} \sum_n \sum_{\mu,\nu,\rho,\sigma} \varepsilon_{\mu\nu\rho\sigma} \int_{f(n,\mu)} d^3x \operatorname{tr} \left[(v_{n,\mu}^{-1} \partial_\nu v_{n,\mu}) (v_{n,\mu}^{-1} \partial_\rho v_{n,\mu}) (v_{n,\mu}^{-1} \partial_\sigma v_{n,\mu}) \right] \\ - \frac{1}{8\pi^2} \sum_n \sum_{\mu,\nu,\rho,\sigma} \varepsilon_{\mu\nu\rho\sigma} \int_{p(n,\mu,\nu)} d^2x \operatorname{tr} \left[(v_{n,\mu} \partial_\rho v_{n,\mu}^{-1}) (v_{n-\hat{\mu},\nu}^{-1} \partial_\sigma v_{n-\hat{\mu},\nu}) \right].$$

- $v_{n,\mu}(x)$ is the gauge translation function (**transition function**).
- On the lattice, topological values are ill-defined.
 - Restricting the size of plaquette (**Admissibility condition**), Lüscher constructed **integral** TC on the lattice (Lüscher, Commun. Math. Phys. 85 (1982)).
 - We aim to construct the **fractional** TC on the $SU(N)$ lattice by extended the Lüscher's topological charge.
 - ✓ Itou, arXiv:1811.05708[hep-th]
 - ✓ Anosova, Gattringer, Göschl, Sulejmanpasic, Törek, arXiv:1912.11685 [hep-lat]

Fiber Bundle


- The fiber bundle describes the gauge theory.
 - Covering a manifold M by patches U_i , each patch has the gauge field a_i and the matter field ϕ_i with the irreducible representation ρ .
- Gauge fields at $U_{ij} = U_i \cap U_j$ are connected by the gauge transformation function g_{ij} .
- At $U_{ijk} = U_i \cap U_j \cap U_k$, the cocycle condition is satisfied,

$$a_j = g_{ij}^{-1} a_i g_{ij} - i g_{ij}^{-1} d g_{ij},$$
$$\phi_j = \rho(g_{ij}^{-1}) \phi_i.$$




transition function g_{ij}

$$g_{ij} g_{jk} g_{ki} = 1.$$



Fiber Bundle on the Lattice

- The manifold is divided by hyper cubes $c(n)$.
- e. g., in the 3d,



Review of Lüscher's construction

Lüscher, Commun. Math. Phys. 85 (1982)

- For the integral topological charge,

Step1


- Define $v_{n,\mu}(n)$ at the corner of $f(n,\mu)$ with complete axial gauge.
➤ Define the parallel transporter $w^m(x)$ with complete axial gauge.

Step2

- Interpolate $v_{n,\mu}(n)$ to the x in the face $f(n,\mu)$.
➤ Define the interpolate function $S_{n,\mu}^m(x)$.

Step3

- To define $S_{n,\mu}^m(x)$ correctly, we need the admissibility condition.




Step1: Transition Function

- Define the parallel transporter ($n \rightarrow x = n + \sum_{\mu} z_{\mu} \hat{\mu}$, $z_{\mu} = \{0,1\}$),

$$w^n(x) := U(n, 4)^{z_4} U(n + z_4 \hat{4}, 3)^{z_3} U(n + z_4 \hat{4} + z_3 \hat{3}, 2)^{z_2} U(n + z_4 \hat{4} + z_3 \hat{3} + z_2 \hat{2}, 1)^{z_1}.$$

- This selection means to take the complete axial gauge on the lattice.
- Gauge transformation on the lattice,

$$U(n, \mu) \mapsto g_n^{-1} U(n, \mu) g_{n+\hat{\mu}}.$$



Step1: Transition Function

- Define the parallel transporter ($n \rightarrow x = n + \sum_{\mu} z_{\mu} \hat{\mu}$, $z_{\mu} = \{0,1\}$),

$$w^n(x) := U(n, 4)^{z_4} U(n + z_4 \hat{4}, 3)^{z_3} U(n + z_4 \hat{4} + z_3 \hat{3}, 2)^{z_2} U(n + z_4 \hat{4} + z_3 \hat{3} + z_2 \hat{2}, 1)^{z_1}.$$

- This selection means to take the complete axial gauge on the lattice.


- Define link variables in complete axial gauge ($n \rightarrow x \rightarrow x + \hat{\mu} \rightarrow n$),

$$u_{x, x+\hat{\mu}}^n := w^n(x) U(x, \mu) w^n(x + \hat{\mu})^{-1}.$$

- Define link variables in complete axial gauge in another way,

$$u_{x, x+\hat{\mu}}^{n-\hat{\mu}} = w^{n-\hat{\mu}}(x) \underbrace{w^n(x)^{-1} u_{x, x+\hat{\mu}}^n w^n(x + \hat{\mu})}_{\sim U(x, \mu)} w^{n-\hat{\mu}}(x + \hat{\mu})^{-1}$$

$$= v_{n, \mu}(x) u_{x, x+\hat{\mu}}^n v_{n, \mu}(x + \hat{\mu})^{-1}.$$



Transition function

$$v_{n, \mu}(x) := w^{n-\hat{\mu}}(x) w^n(x)^{-1}$$

Step2: To the Coordinate x

- Interpolate the transition function to the x in the face $f(n, \mu)$,

$$v_{n,\mu}(x) = S_{n,\mu}^{n-\hat{\mu}}(x)^{-1} v_{n,\mu}(n) S_{n,\mu}^n(x).$$

- Define the interpolate function $S_{n,\mu}^m(x)$.

$$f_{n,\mu}^m(x_\gamma) = (u_{s_3 s_0}^m)^{y_\gamma} (u_{s_0 s_3}^m u_{s_3 s_7}^m u_{s_7 s_2}^m u_{s_2 s_0}^m)^{y_\gamma} u_{s_0 s_2}^m (u_{s_2 s_7}^m)^{y_\gamma},$$



$$g_{n,\mu}^m(x_\gamma) = (u_{s_5 s_1}^m)^{y_\gamma} (u_{s_1 s_5}^m u_{s_5 s_4}^m u_{s_4 s_6}^m u_{s_6 s_1}^m)^{y_\gamma} u_{s_1 s_6}^m (u_{s_6 s_4}^m)^{y_\gamma},$$

$$h_{n,\mu}^m(x_\gamma) = (u_{s_3 s_0}^m)^{y_\gamma} (u_{s_0 s_3}^m u_{s_3 s_5}^m u_{s_5 s_1}^m u_{s_1 s_0}^m)^{y_\gamma} u_{s_0 s_1}^m (u_{s_1 5}^m)^{y_\gamma},$$

$$k_{n,\mu}^m(x_\gamma) = (u_{s_7 s_2}^m)^{y_\gamma} (u_{s_2 s_7}^m u_{s_7 s_4}^m u_{s_4 s_6}^m u_{s_6 s_2}^m)^{y_\gamma} u_{s_2 s_6}^m (u_{6 s_4}^m)^{y_\gamma},$$

$$l_{n,\mu}^m(x_\beta, x_\gamma) = [f_{n,\mu}^m(x_\gamma)^{-1}]^{y_\beta} [f_{n,\mu}^m(x_\gamma) k_{n,\mu}^m(x_\gamma) g_{n,\mu}^m(x_\gamma)^{-1} h_{n,\mu}^m(x_\gamma)^{-1}]^{y_\beta} \cdot h_{n,\mu}^m(x_\gamma) [g_{n,\mu}^m(x_\gamma)]^{y_\beta},$$

$$S_{n,\mu}^m(x_\alpha, x_\beta, x_\gamma) = (u_{s_0 s_3}^m)^{y_\gamma} [f_{n,\mu}^m(x_\gamma)]^{y_\beta} [l_{n,\mu}^m(x_\beta, x_\gamma)]^{y_\alpha}.$$



Step2: To the Coordinate x

- Interpolate the transition function to the x in the face $f(n, \mu)$,

$$v_{n,\mu}(x) = S_{n,\mu}^{n-\hat{\mu}}(x)^{-1} v_{n,\mu}(n) S_{n,\mu}^n(x).$$



- e.g., in 2d, ($0 \leq y_1 \leq 1$)

$$S_{n,2}^{n-\hat{2}}(x) = U(n - \hat{2}, 1)^{y_1} U(n - \hat{2} + \hat{1}, 2)^{y_1} U(n - \hat{2}, 2)^{-y_1},$$

$$S_{n,2}^n(x) = U(n, 1)^{y_1},$$

$$v_{n,2}(n) = U(n - \hat{2}, 2),$$

$$\begin{aligned} v_{n,2}(x) &= S_{n,2}^{n-\hat{2}}(x)^{-1} v_{n,2}(n) S_{n,2}^n(x) \\ &= U(n - \hat{2}, 2) \exp [iy_1 F_{12}(n - \hat{2})]. \end{aligned}$$



Step3: Admissibility condition

- We take the compact $SU(N)$ lattice gauge theory.
- For simplicity, in the compact $U(1)$ lattice gauge theory,

$$U_\mu(n) = e^{ia_\mu(n)}, \quad (-\pi \leq a_\mu(n) \leq \pi)$$

$$F_{\mu\nu}(n) = \frac{1}{i} \ln U_\mu(n)U_\nu(n + \hat{\mu})U_\mu(n + \hat{\nu})^{-1}U_\nu(n)^{-1} := \frac{1}{i} \ln U_p. \quad (-\pi \leq F_{\mu\nu}(n) \leq \pi)$$

- When $|F_{\mu\nu}| = \pi$, the component, $e^{iyF_{\mu\nu}} = (-1)^y$, becomes ambiguous.
- Require the gauge field smooth (**admissibility condition**),

$$\|\mathbb{1} - U_p\| < \varepsilon.$$


Step3: Admissibility condition

- We take the compact $SU(N)$ lattice gauge theory.
- Considering the interpolate function, we select the value of ε .

$$\|\mathbb{1} - U_p\| < \varepsilon.$$

- Recall the link variables with the complete axial gauge and interpolate function.

➤ e.g.,



In the component of the interpolate function, plaquette is appeared.

$$\begin{aligned}
 f_{n,\mu}^m(x_\gamma) &= (u_{s_3 s_0}^m)^{y_\gamma} (u_{s_0 s_3}^m u_{s_3 s_7}^m u_{s_7 s_2}^m u_{s_2 s_0}^m)^{y_\gamma} u_{s_0 s_2}^m (u_{s_2 s_7}^m)^{y_\gamma}, \\
 g_{n,\mu}^m(x_\gamma) &= (u_{s_5 s_1}^m)^{y_\gamma} (u_{s_1 s_5}^m u_{s_5 s_4}^m u_{s_4 s_6}^m u_{s_6 s_1}^m)^{y_\gamma} u_{s_1 s_6}^m (u_{s_6 s_4}^m)^{y_\gamma}, \\
 h_{n,\mu}^m(x_\gamma) &= (u_{s_3 s_0}^m)^{y_\gamma} (u_{s_0 s_3}^m u_{s_3 s_5}^m u_{s_5 s_1}^m u_{s_1 s_0}^m)^{y_\gamma} u_{s_0 s_1}^m (u_{s_1 s_5}^m)^{y_\gamma}, \\
 k_{n,\mu}^m(x_\gamma) &= (u_{s_7 s_2}^m)^{y_\gamma} (u_{s_2 s_7}^m u_{s_7 s_4}^m u_{s_4 s_6}^m u_{s_6 s_2}^m)^{y_\gamma} u_{s_2 s_6}^m (u_{s_6 s_4}^m)^{y_\gamma}, \\
 l_{n,\mu}^m(x_\beta, x_\gamma) &= [f_{n,\mu}^m(x_\gamma)^{-1}]^{y_\beta} [f_{n,\mu}^m(x_\gamma) k_{n,\mu}^m(x_\gamma) g_{n,\mu}^m(x_\gamma)^{-1} h_{n,\mu}^m(x_\gamma)^{-1}]^{y_\beta} \\
 &\quad \cdot h_{n,\mu}^m(x_\gamma) [g_{n,\mu}^m(x_\gamma)]^{y_\beta}, \\
 S_{n,\mu}^m(x_\alpha, x_\beta, x_\gamma) &= (u_{s_0 s_3}^m)^{y_\gamma} [f_{n,\mu}^m(x_\gamma)]^{y_\beta} [l_{n,\mu}^m(x_\beta, x_\gamma)]^{y_\alpha}.
 \end{aligned}$$

Summary of Lüscher's construction

- Define the transition function at the coordinate $x \in f(n, \mu)$,

$$v_{n,\mu}(x) = S_{n,\mu}^{n-\hat{\mu}}(x)^{-1} v_{n,\mu}(n) S_{n,\mu}^n(x).$$


- ✓ Satisfy the cocycle condition,

$$v_{n-\hat{\mu},\nu}(x) v_{n,\mu}(x) v_{n,\nu}(x)^{-1} v_{n-\hat{\nu},\mu}(x)^{-1} = \mathbb{1}.$$

- Substituting $v_{n,\mu}(x)$, calculate integral topological charge.
- Define the admissibility condition,

$$\|\mathbb{1} - U_p\| < \varepsilon.$$

- Extend these to the fractional topological charge.



\mathbb{Z}_N 1-form Global Transformation on the Lattice

- Lattice $SU(N)$ gauge theory, the action is

$$S_W[U_l] \equiv \sum_p \beta [\text{tr}(\mathbb{1} - U_p) + \text{c.c.}].$$

- Center transformation (\mathbb{Z}_N 1-form global transformation) on the lattice acts on the link variables,


$$U_l \mapsto e^{\frac{2\pi i}{N}k} U_l, \quad W(C) \mapsto e^{\frac{2\pi i}{N}k} W(C).$$

- ✂ Recall that under the 1-form global transformation in the continuum theory, the Wilson line changes,

$$\langle U_\alpha(\Sigma) W(C) \dots \rangle = e^{-i\alpha} \langle W(C) \dots \rangle$$

- The transition function satisfies the cocycle condition still.

$$v_{n-\hat{\mu},\nu}(x) v_{n,\mu}(x) v_{n,\nu}(x)^{-1} v_{n-\hat{\nu},\mu}(x)^{-1} = \mathbb{1}.$$



\mathbb{Z}_N 1-form Gauge Transformation on the Lattice

- Gauging the center symmetry, the action becomes

$$S_W[U_\ell, B_p] = \sum_p \beta \left[\text{tr} \left(\mathbb{1} - e^{-\frac{2\pi i}{N} B_p U_p} \right) + \text{c.c.} \right].$$

- Invariant under the \mathbb{Z}_N 1-form gauge transformation,

$$U_\ell \mapsto e^{\frac{2\pi i}{N} \lambda_\ell} U_\ell, \quad B_p \mapsto B_p + (d\lambda)_p.$$


- The cocycle condition is relaxed,

$$\tilde{v}_{n-\hat{\nu}, \mu}(n) \tilde{v}_{n, \nu}(n) \tilde{v}_{n, \mu}(n)^{-1} \tilde{v}_{n-\hat{\mu}, \nu}(n)^{-1} = e^{\frac{2\pi i}{N} B_{\mu\nu}(n-\hat{\mu}-\hat{\nu})} \mathbb{1}.$$

- ✂ 't Hooft twisted boundary condition

$$U(n + L\hat{\nu}, \mu) = g_{n, \mu}^{-1} U(n, \mu) g_{n+\hat{\mu}, \nu}$$

$$g_{n+L\hat{\nu}, \mu}^{-1} g_{n, \nu}^{-1} g_{n, \mu} g_{n+L\hat{\mu}, \nu} = e^{\frac{2\pi i}{N} z_{\mu\nu}}, \quad z_{\mu\nu} = \sum_p B_p \text{ mod } N.$$



Transition Function for Fractional TC

- Coupling \mathbb{Z}_N 2-form field to the theory, the structure of fiber bundle becomes rich.

$$\tilde{v}_{n-\hat{\nu},\mu}(n)\tilde{v}_{n,\nu}(n)\tilde{v}_{n,\mu}(n)^{-1}\tilde{v}_{n-\hat{\mu},\nu}(n)^{-1} = e^{\frac{2\pi i}{N}B_{\mu\nu}(n-\hat{\mu}-\hat{\nu})}\mathbb{1}.$$

- We find that **the \mathbb{Z}_N 1-form gauge invariance** plays the center role.


➤ Admissibility condition

$$\|\mathbb{1} - \tilde{U}_{\mu\nu}(n)\| < \varepsilon,$$

$$\tilde{U}_{\mu\nu}(n) \equiv e^{-\frac{2\pi i}{N}B_{\mu\nu}(n)}$$

$$\times U(n, \mu)U(n + \hat{\mu}, \nu)U(n + \hat{\nu}, \mu)^{-1}U(n, \nu)^{-1}.$$

➤ Components of transition function



Construction of $u_{s_7 s_2}^{n-3}$

Construction of $\tilde{u}_{s_7 s_2}^{n-3}$

Fractional TC

- By the \mathbb{Z}_N 1-form invariant transition function, we calculate TC,

$$z_{\mu\nu} = \sum_{p \in (T^2)_{\mu\nu}} B_p \pmod{N},$$

$$Q_{\text{top}} = -\frac{1}{N} \int_{T^4} \frac{1}{2} P_2(B_p) \pmod{1} \in -\frac{1}{N} \frac{\varepsilon_{\mu\nu\rho\sigma} z_{\mu\nu} z_{\rho\sigma}}{8} + \mathbb{Z},$$

$$P_2(B_p) = B_p \cup B_p + B_p \cup_1 dB_p.$$

- In the $U(1)$ lattice gauge theory, we make sure that (cf. Abe, Morikawa, Suzuki, arXiv:2210.12967[hep-th])

$$Q_{\text{top}} = \frac{1}{32\pi^2} \sum_{n \in \Lambda} \sum_{\mu, \nu, \rho, \sigma} \varepsilon_{\mu\nu\rho\sigma} \tilde{F}_{\mu\nu}(n) \tilde{F}_{\rho\sigma}(n + \hat{\mu} + \hat{\nu}) \in \frac{1}{N^2} \mathbb{Z} + \mathbb{Z}.$$

Anomaly I

- Again, the action on the lattice is

$$S[U_l, B_p] \equiv -S_W[U_l, B_p] + i\theta Q_{\text{top}}[U_l, B_p].$$

- The topological charge is

$$Q_{\text{top}} = -\frac{1}{N} \int_{T^4} \frac{1}{2} P_2(B_p) + \mathbb{Z} \equiv \text{frac}[B_p] + \text{int}[U_l, B_p].$$

✧ Manifestly invariant under the \mathbb{Z}_N one-form gauge transformation

- We discuss the anomaly at $\theta = \pi$ between the \mathbb{Z}_N 1-form gauge and θ shift.

Anomaly II

- At $\theta = \pi$, the partition function is, under \mathcal{T} transformation,

$$Z[B_p] = \int \mathcal{D}U_l e^{S[U_l, B_p]} = \int \mathcal{D}U_l e^{-S_W[U_l, B_p]} e^{i\theta Q_{\text{top}}[U_l, B_p]}$$

$$\begin{aligned} \xrightarrow{\theta=\pi, \theta \text{ shift.}} Z'[B_p] &= \int \mathcal{D}U_l e^{-S_W[U_l, B_p]} e^{i\pi(-Q_{\text{top}}[U_l, B_p])} = \int \mathcal{D}U_l e^{-S_W[U_l, B_p]} e^{i\pi Q_{\text{top}}[U_l, B_p]} \underbrace{e^{-i2\pi Q_{\text{top}}[U_l, B_p]}}_{=e^{-i2\pi \text{int}[U_l, B_p]} e^{-i2\pi \text{frac}[B_p]}} \\ &= e^{-i2\pi \text{frac}[B_p]} \underbrace{\int \mathcal{D}U_l e^{-S_W[U_l, B_p]} e^{i\pi Q_{\text{top}}[U_l, B_p]}}_{=Z} \neq Z[B_p] \end{aligned}$$

- This means that there is the anomaly at $\theta = \pi$ between the \mathbb{Z}_N 1-form gauge and θ shift.

Conclusion and Future Work

☆ Conclusion

- We construct the fractional topological charge on the $SU(N)$ lattice gauge theory.
- By this topological charge, we construct the anomaly at $\theta = \pi$ between the \mathbb{Z}_N 1-form gauge and θ shift on the lattice.

☆ Future work

- Construct the magnetic operator under the admissibility condition on the lattice
 - ✓ cf. Abe, Morikawa, Onoda, Suzuki, Tanizaki, arXiv:2304.14815 [hep-lat]
- Construct non-invertible symmetries on the lattice